

## **Abstracts of the section ‘Innovative and Applied Research in Biology’ at the 83rd International Scientific Conference of the University of Latvia**

**The International Scientific Conference of the University of Latvia** has become an integral part of scientific life, bringing along nearly centuries of tradition. This year, the 83rd International Scientific Conference of the University of Latvia took place from January to March 2025. The conference sessions were organised into seven thematic sections: Exact Sciences and Technology, Education Sciences and Psychology, Law, Medicine and Life Sciences, Economics, Social Sciences, Humanities, and Interdisciplinary Studies. Each section encompassed a wide range of topics, covering significant research directions and innovation fields. More than 2500 speakers – students, faculty academic staff, researchers, as well as representatives of various companies and organisations – participated in section meetings organised by faculties, the Botanical Garden, the University of Latvia Museum, the University of Latvia Library, and regional branches. Remote working opportunities had facilitated the broad participation of foreign researchers from various continents and scientific institutions, as well as attracted more listeners, allowing a wider audience to discuss the latest scientific advances. Participating in the conference provided an excellent opportunity to share your research results, introduced the latest scientific achievements, and established connections with colleagues in Latvia and other countries.

On 6 March 2025, the section **‘Innovative and Applied Research in Biology’** was organised for the seventh time in Riga by the Institute of Biology. The main focus of this conference section was on innovative and applied research in biology and interdisciplinary fields. The International Scientific Committee included Dr biol. Liene Aunina (LU, Latvia), Dr biol. Mārtiņš Briedis (Swiss Ornithological Institute, Switzerland), Dr biol. Dalius Butkauskas (Nature Research Centre, Lithuania), Dr biol. Dmitry Telnov (Natural History Museum, London, UK), Dr biol. Dace Grauda (LU, Latvia) Dr biol. Anita Osvalde (LU, Latvia), Dr biol. Pāvels Semjonovs (LU, Latvia), and Dr biol. Īzaks Rašāls (LU, Latvia)

Compiler: Dr biol. Līga Jankevica

Editors: Līga Jankevica and Dmitry Telnov

**Oral presentations****O1: APPLICATION OF MOLECULAR METHODS IN ZOOLOGY IN THE BALTIC COUNTRIES****Algimantas Paulauskas\****Vytautas Magnus University, Kaunas, Lithuania**\* Corresponding author. Email: algimantas.paulauskas@vdu.lt*

---

Contemporary tools of molecular biology open new possibilities in zoology and provide important answers to classic problems. Zoological questions of mating strategies, physiological adaptation, identification of life-cycle stages, genetic exchange between populations, analysis of hybridisation, population structure in 'typical' sexual organisms, measuring inbreeding, species/subspecies identification (a technique for species identification of parasites without the need for separation from the host), determination of geographical origin, and many others are now being powerfully addressed using tools from the molecular arsenal. Molecular genetic techniques are clearly having an impact on what has been traditionally considered to be purely zoological studies. Applications in the Baltic countries are discussed.

**Keywords:** molecular methods, zoology, application

**References**

1. Baverstock PR. Applications of molecular genetic techniques in zoology. *Zool Rev.* 1988;1:1–13.
  2. Molecular zoology advances, strategies and protocols. Edited by Ferraris JD and Palumbi SR. New York: J. Wiley; 1996. 580 p.
  3. Chambers GK, Curtis C, Millar CD, et al. DNA fingerprinting in zoology: past, present, future. *Investig Genet.* 2014;5:3.
  4. International Journal of Molecular Zoology (ISSN 1927-534X).
- 

**O2: DETECTION AND PHYLOGENETIC CHARACTERISATION OF TICK-BORNE ENCEPHALITIS VIRUS IN LITHUANIA****Jana Radzijeuskaja\*, Algimantas Paulauskas***Vytautas Magnus University, Kaunas, Lithuania**\* Corresponding author. Email: jana.radzijeuskaja@vdu.lt*

---

The Baltic countries are the region in Europe where tick-borne encephalitis (TBE) is most prevalent. Lithuania has the highest notification rate of TBE cases, with a notable increase in incidence since 1992. A recent study reported a 0.4% prevalence of the TBE virus (TBEV) in Lithuania's two most common tick species, *Ixodes ricinus* and *Dermacentor reticulatus*. TBEV was detected in the three developmental stages (adults, nymphs, and larvae) of *I. ricinus* and *D. reticulatus* adults. TBEV-infected ticks were found in 16 locations in seven counties, with

prevalence ranging from 0.1% to 1.0%. In Lithuania, TBEV was also detected in different rodent species, different breeds of dogs, horses, and goat and sheep milk samples. While TBE is a significant health issue in Lithuania, detailed information about local virus strains and their genetic variability is still limited. The present study compared nucleotide sequences of the E and NS3 genes of 39 TBEV strains isolated from ticks, rodents, and humans in Lithuania over a period of 20 years. Phylogenetic analysis of the partial E and NS3 gene sequences from the TBEV isolates indicated that these strains were specific to Lithuania and belonged to the European subtype, exhibiting significant regional genetic diversity. The analysis showed that certain TBEV strains were specific to particular regions. In most cases, TBEV strains from the same sampling site were identical; however, genetic differences in the E and NS3 gene sequences were noted among strains from different geographical areas of Lithuania. The detected TBEV genotypes were not specific to the tick species. TBEV has a focal distributional pattern in endemic areas in Lithuania as the virus is not uniformly present in the tick population. The present study indicates that at least six distinct virus lineages are circulating among ticks and two among rodents in the country. Further research involving whole-genome sequencing is necessary to enhance the understanding of the regional genetic diversity of TBEV strains.

**Keywords:** tick-borne encephalitis, TBE virus, genetic diversity, Lithuania

#### References

1. Sidorenko M, Radzijeuskaja J, Mickevičius S, Bratčikovienė N, Paulauskas A. Prevalence of tick-borne encephalitis virus in questing *Dermacentor reticulatus* and *Ixodes ricinus* ticks in Lithuania. *Ticks Tick Borne Dis.* 2021;12:101594.
2. Sidorenko M, Radzijeuskaja J, Mickevičius S, Bratčikov M, Mardosaitė-Busaitienė D, Sakalauskas P, Paulauskas A. Phylogenetic characterisation of tick-borne encephalitis virus from Lithuania. *PLoS One.* 2024;19:e0296472.
3. Simkute E, Pautienius A, Grigas J, Sidorenko M, Radzijeuskaja J, Paulauskas A, Stankevicius A. The prevalence of tick-borne encephalitis virus in wild rodents captured in tick-borne encephalitis foci in highly endemic Lithuania. *Viruses.* 2024;16:444.

---

## O3: GENETIC VARIABILITY OF AMERICAN MINK (*NEOVISON VISON*) IN LITHUANIA

Daiva Šakienė, Asta Aleksandravičienė, Loreta Gričiuvienė, Algimantas Paulauskas\*

*Vytautas Magnus University, Kaunas, Lithuania*

\* Corresponding author. Email: algimantas.paulauskas@vdu.lt

---

The American mink (*Neovison vison*), a species native to North America, has been successfully introduced and established in the wild across Europe. As a result, it has largely displaced the native European mink (*Mustela lutreola*). In Lithuania, the American mink has also been observed to negatively impact native wildlife, including small mammals, fish, and amphibians, while contributing to broader ecological imbalances and disrupting local biodiversity. The aim of this study was to assess the genetic variability of different American mink (*Neovison vison*) subpopulations in Lithuania. The tissue samples of *N. vison* were collected from three Lithuania localities: Alytus, Šilutė, and Zarasai. Genomic DNA was extracted from tissue samples using the Genomic DNA Purification Kit (Thermo Fisher Scientific, Lithuania) following the manufacturer's instructions. To assess the genetic diversity of the American mink, six microsatellite markers were used. The results showed that, within each subpopulation, observed heterozygosity was higher than expected

heterozygosity, suggesting possible factors such as genetic admixture, or population bottlenecks. Among the populations studied, the Alytus subpopulation displayed the lowest genetic diversity, while the Šilutė subpopulation showed the highest. Principal Coordinates Analysis (PCoA) revealed notable genetic differentiation between Alytus and Zarasai subpopulations, highlighting significant genetic distances between these geographically separated groups. Additionally, the genetic structure of the Šilutė subpopulation showed similarities to both Alytus and Zarasai subpopulations. This pattern suggests gene flow or shared ancestry, despite the geographical separation by different river basins (the Nemunas River and the Dauguva River basins, respectively). These results provide valuable insights into the genetic diversity and population structure of the introduced *N. vison* in Lithuania, which may inform future management and conservation strategies aimed at mitigating their ecological impact on native species.

**Keywords:** Lithuania, genetic variability, *Neovison vison*, microsatellite markers

#### References

1. Lecis R, Ferrando A, Ruiz-Olmo I, Manas S, Domingo-Roura X. Population genetic structure and distribution of introduced American mink (*Mustela vison*) in Spain, based on microsatellite variation. *Conserv Genet.* 2008;9:1149–61.
2. Michalska-Parda A, Brzeziński M, Zalewski A, Kozakiewicz M. Genetic variability of feral and ranch American mink *Neovison vison* in Poland. *Acta Theriologica.* 2009;54:1–10.

---

## O4: 3D BIOTEXTILE MANUFACTURING AND ASSESSMENT OF PROTECTIVE PROPERTIES

**Dalius Butkauskas<sup>1\*</sup>, Dace Grauda<sup>2</sup>, Inga Lashenko<sup>3</sup>**

<sup>1</sup> *Nature Research Centre, Vilnius, Lithuania*

<sup>2</sup> *Institute of Biology, FMLS, University of Latvia, Riga, Latvia*

<sup>3</sup> *JLU Technologies Ltd, Riga, Latvia*

\* *Corresponding author. Email: dalius.butkauskas@gamtc.lt*

---

The development of 3D textile models for biological testing began with the production of biotextile material by the project partner A Grupe (Jonava, Lithuania), a company specialising in linen fabrics and products. To enhance the protective properties of these newly created biotextiles, which consisted of high-quality, 3D fibre composite variants, the materials were impregnated with an aqueous solution. This solution contained 0.5% polyethylene oxide, 0.5% hydroxyethyl cellulose, and 0.5–1% silicone. The resulting polymer structures coating the fabrics also incorporated succinic acid particles (or succinic acid derivatives) in some versions, along with SiO<sub>2</sub> and Ag nanoparticles in others, creating a range of different polymer matrices. The impregnation procedures and production of the various biotextile versions were carried out at JLU Technologies (Riga, Latvia). Biological testing, focused on the protective properties of the biotextile materials, was conducted at the Laboratory of Environmental Genetics of the FMLS Institute of Biology University of Latvia and the Laboratory of Molecular Ecology of the Nature Research Centre (Lithuania). Common duckweed (*Lemna minor*) and fruit flies (*Drosophila melanogaster*) were chosen as the primary test organisms due to their cost-effectiveness, reproducibility, and reliability in experiments.

To investigate the protective properties of the biotextiles, we examined the effects of low-frequency electromagnetic radiation (LF EMF, 50 Hz, 1.3 mT magnetic flux density) on the physiological and molecular levels of the test organisms. *L. minor* (diploid laboratory line Sta2 axenic



culture) plants were grown in Petri dishes wrapped in different biotextile versions and placed within a Helmholtz coil. The growth rate, the number of fronds, and the total frond area increase per week were measured. After one week, DNA was collected to assess the potential of LF EMF to induce point mutations in genes involved in oxidative stress regulation. Nuclear ascorbate peroxidase (APx) and catalase (Cat) gene fragments, along with selected chloroplast microsatellite loci, were sequenced.

Results showed a reduction in point mutations in *L. minor* Sta2 clones grown with biotextiles containing 0.25%, 0.5%, and 1% succinite nanoparticles compared to the control group (biotextile without succinite). Conversely, biotextiles containing 0.5% succinic acid or Ag nanoparticles showed no positive effect on mutation reduction and significantly suppressed growth intensity, leading to increased point mutations in the APx and Cat gene fragments and microsatellite loci, and decreased frond area.

The positive effect of succinite nanoparticles, particularly at low concentrations, in mitigating point mutations contrasted with the opposite effect of succinic acid. The biotextile containing 0.5% succinite, 0.1% SiO<sub>2</sub>, and 0.1% Ag nanoparticles highlighted the importance of nanoparticle origin and concentration, as well as the sensitivity of the *L. minor* Sta2 model, demonstrating its potential for biotesting applications.

In conclusion, 3D biotextile versions incorporating succinite nanoparticles (0.25% to 1%) within the polymer fabric coating demonstrate significant potential as protective biotextiles, mitigating the effects of LF EMF and potentially other stressors.

**Keywords:** succinite nanoparticles, succinic acid, SiO<sub>2</sub>, Ag nanoparticles, *L. minor*, low frequency electromagnetic field 50 Hz, ascorbate peroxidase, catalase, microsatellite loci, point mutations

**Acknowledgements:** the study was financially supported by the S-M-ERA.NET-22-1 project 3DNano-HPC.

---

## O5: IMPACT OF AMBER NANOPARTICLES AND ITS DERIVATIVES ON *L. MINOR* CLONE AFFECTED BY LOW FREQUENCY ELECTROMAGNETIC RADIATION (50 Hz)

Ieva Ignatavičienė<sup>\*1</sup>, Regina Vyšniauskienė<sup>1</sup>, Vida Rancėlienė<sup>1</sup>, Rimantas Petrošius<sup>1</sup>,  
Dace Grauda<sup>2</sup>, Dalius Butkauskas<sup>1</sup>

<sup>1</sup> Nature Research Centre, Vilnius, Lithuania

<sup>2</sup> Institute of Biology, University of Latvia, Riga, Latvia

\* Corresponding author. Email: ieva.ignataviciene@gamtc.lt

---

Textile technologies have recently attracted great attention as potential tools for biomedical application or protection from stressful environments (Rivero et al., 2015). Some natural organic compounds with fungicidal and antibacterial properties can be included in the silicon structure covering biotextile materials to enhance their functional properties. Although it is known that amber or its derivatives have healing properties (Rissanen et al., 2021), the effect on living organisms remains unanswered. In this work, the effect of 3D biotextile with incorporated 0.25%, 0.5%, and 1% concentration of amber, succinic acid, succinic acid with SiO<sub>2</sub> and combining 0.5% amber, 0.1% SiO<sub>2</sub> and 0.1% Ag nanoparticles were investigated. In the study, plants of *L. minor* laboratory clone Sta2 grown in a Petri dishes wrapped within 3D bio-textile specimens embedded with

different nanoparticles were exposed to 50 Hz electromagnetic radiation for seven days. After the termination of the experiment, the frond area of *L. minor* plants was examined molecularly by sequencing nuclear ascorbate peroxidase (APX), catalase (CAT) gene fragments, and some chloroplast microsatellite loci. In most cases a reduction of point mutations and an increment of frond area was detected in *L. minor* clones grown coated with bio-textile encompassing 0.25%, 0.5%, and 1% amber nanoparticles compared to control group that was grown coated with the same bio-textile material that doesn't contain amber particles. The opposite effect was observed by adding 0.5% succinic acid or 0.1% Ag nanoparticles into 3D bio-textile, as the number of point mutations significantly increased APX, CAT gene fragments and microsatellite loci, and the frond area decreased in comparison to the control group. Our results demonstrate a positive effect of amber nanoparticles, especially at low concentrations of amber, of promoting growth and lowering the number of point mutations. The opposite negative effect of succinic acid alone and composition of 0.5% amber, 0.1% SiO<sub>2</sub> and 0.1% Ag incorporated into the biotextile, enhanced the number of point mutations and decreased the growth of the *L. minor* clones exposed to electromagnetic radiation. The 3D biotextile containing amber nanoparticles have significant potential to be used as a protective tissue in stressful conditions.

**Keywords:** 3D bio-textile, amber nanoparticle, succinic acid nanoparticle, silver nanoparticle, *L. minor*, 50 Hz electromagnetic radiation

**Acknowledgements:** the study was financially supported by the S-MERA. NET-22-1 project 3DNano-HPC.

#### References

1. Rivero PJ, Urrutia A, Goicoechea J, Arregui FJ. Nanomaterials for functional textiles and fibers. *Nanoscale Res Lett.* 2015;501:10.
2. Rissanen K, Hölttä T, Bäck J, Rigling A, Wermelinger B, Gessler A. Drought effects on carbon allocation to resin defences and on resin dynamics in old-grown Scots pine. *Environ Exp Bot.* 2011;185:104410.

---

## O6: ANTIMICROBIAL EFFICACY OF DIFFERENT *ROSA RUGOSA* PARTS EXTRACTED USING AQUEOUS AND ETHANOLIC EXTRACT

Žaneta Maželienė<sup>1,3\*</sup>, Jolita Kirvaitienė<sup>1,2</sup>, Giedrė Jarienė<sup>1</sup>, Asta Aleksandravičienė<sup>1,4</sup>, Rasa Volskienė<sup>1</sup>

<sup>1</sup> Faculty of Medicine, Kauno Kolegija Higher Education Institution, Kaunas, Lithuania

<sup>2</sup> Faculty of Public Health, Lithuanian University of Health Sciences, Kaunas, Lithuania

<sup>3</sup> Faculty of Veterinary Medicine, Institute of Microbiology and Virology, Lithuanian University of Health Sciences, Kaunas, Lithuania

<sup>4</sup> Faculty of Natural Sciences, Vytautas Magnus University, Kaunas, Lithuania

\* Corresponding author. Email: zaneta.mazeliene@go.kauko.lt

---

With the increasing prevalence of drug-resistant pathogens, the search for alternative antimicrobial agents has intensified. Bioactive compounds derived from medicinal plants have gained significant attention as they offer a promising natural source of novel antifungal and antibacterial agents with potentially fewer side effects compared to synthetic drugs. *Rosa rugosa* has been traditionally used in herbal medicine for its diverse therapeutic properties.

The aim of this study was to evaluate the antimicrobial potential of aqueous and ethanolic extracts obtained from various parts of *R. rugosa*, including rose hips, flowers, petals, leaves, stems, and roots. The antimicrobial efficacy of the extracts was assessed using agar well diffusion method under controlled laboratory conditions. To ensure extract purity and enhance the concentration of active compounds, a rotary evaporator was employed for solvent removal. The antimicrobial activity was tested against a panel of microbial strains, including *Candida albicans* (a common fungal pathogen), four gram-positive bacterial strains (*Staphylococcus aureus*, *Bacillus subtilis*, *Enterococcus faecalis*, and *Listeria monocytogenes*), and four gram-negative bacterial strains (*Escherichia coli*, *Pseudomonas aeruginosa*, *Salmonella enterica*, and *Klebsiella pneumoniae*). The results indicated that extracts from different plant parts exhibited varying degrees of antimicrobial activity. *R. rugosa* leaf extracts demonstrated the highest antimicrobial effectiveness, followed by petal and rose hip extracts, while root extracts showed moderate activity. In contrast, the weakest antimicrobial effects were observed in extracts derived from stems and rose hips. The study also revealed that ethanolic extracts had superior antimicrobial activity compared to aqueous extracts, highlighting the potential of ethanol as a more effective solvent for extracting bioactive compounds. Both aqueous and ethanolic extracts exhibited notable antifungal activity against *Candida albicans*, with ethanolic extracts proving to be more potent in inhibiting fungal growth. Among all tested plant parts, petals exhibited the strongest antifungal properties, suggesting a high concentration of antifungal compounds in these tissues.

In conclusion, this study provides valuable insights into the antimicrobial potential of *R. rugosa* extracts and underscores their potential use in developing natural antimicrobial agents. Further research is needed to isolate and characterise the specific bioactive compounds responsible for the observed effects and to explore their possible applications in the pharmaceutical and food industries.

**Keywords:** antimicrobial efficacy, *Rosa rugosa*, aqueous extract, ethanolic extract

#### References

1. Cendrowski A, Kraśniewska K, Przybył JL, Zielińska A, Kalisz S. Antibacterial and antioxidant activity of extracts from Rose Fruits (*Rosa rugosa*). *Molecules*. 2020;25:1365.
2. Ribotta S, Liccari F, Muggia L, Pallavicini A, Bagnolini F, Tordoni E, Bacaro G. Invasion at the edge: the case of *Rosa rugosa* (Rosaceae) in Italy. *Diversity*. 2021;13:645.

---

## O7: SHADE TREATMENT EFFECT ON CHLOROPHYLL CONTENT AND GROWTH OF CLODBERRY *RUBUS CHAMAEMORUS*

Laura Āboliņa<sup>1,2\*</sup>, Andis Karlsons<sup>2</sup>, Anita Osvalde<sup>2</sup>

<sup>1</sup> Faculty of Agriculture and Food technology, Latvia University of Life Sciences and Technologies, Jelgava, Latvia

<sup>2</sup> Laboratory of Plant Mineral Nutrition, Institute of Biology, FMLS, University of Latvia, Riga, Latvia

\* Corresponding author. Email: abolina.la@gmail.com

---

Cloudberry is traditionally harvested in the wild in Latvia. Recently, the development of cloudberry cultivation technology has begun. Extracted peat bog recultivation methods include berry cultivation, thus cloudberry could be another perspective berry crop in Latvia. Cloudberries are nutritious and highly demanded for their taste in the northern regions (Tahvanainen et al. 2022), thus cultivation would significantly increase the supply. Netting is a protection method for berry

crops: it decreases high temperatures and provides mechanical protection (Kotilainen et al. 2018). Considering that cloudberry prefer partly shaded environment in the wild, this study was conducted to assess the influence of different net shade levels on cloudberry chlorophyll content and overall development under semi-controlled conditions. Propagated cloudberry were planted outdoors and grown under four shading treatments: control (no shade), 30%, 50%, and 80% shade for two years (2023–2024). SPAD measurements were taken weekly from May to August in both years, while leaf size, density, and nutrient concentrations were determined at the end of August 2024. The highest SPAD values, leaf size, and nutrient contents were observed in plants under 80% shade in both years. In the second year, plants under 50% shade exhibited similar results to those under 80% shade. After two years of growth, the highest plant vitality was observed under 50% and 80% shade, as indicated by plant colour and significantly higher chlorophyll content. However, the shaded plants developed larger leaves to increase light capture, a typical adaptation of shade-tolerant species. Leaves under 80% shade were significantly larger than the control and 30% shade treatments, while the 50% shade treatment was similar to the other treatments. In berry production, increased vegetative growth occurs at the expense of fruit production (Davenport et al. 1996). Therefore, 50% shade is considered the most effective and energy-efficient for cloudberry in the Latvian climate, as indicated by our study. Further studies regarding the combined effects of shade and fertilisation on cloudberry production are required, as optimal nutrient availability is essential for plant development and maximising berry yield.

**Keywords:** netting, SPAD, berry cultivation, light capture

**Acknowledgements:** this research was supported by project ‘Strengthening the Institutional Capacity of LBTU for Excellence in Studies and Research’, grant No. 5.2.1.1.i.0/2/24/I/CFLA/002, funded by the Recovery and Resilience Facility.

#### References

1. Davenport JR. The effect of nitrogen fertilizer rates and timing on cranberry yield and fruit quality. *J Am Soc Hortic Sci.* 1996;121:1089–94.
2. Kotilainen T, Robson TM, Hernández R. Light quality characterization under climate screens and shade nets for controlled-environment agriculture. *PLOS ONE.* 2018;13:e0199628.
3. Tahvanainen V, Miina J, Kurttila M. Climatic and economic factors affecting the annual supply of wild edible mushrooms and berries in Finland. *Forests.* 2019;10:385.

---

## O8: GROWTH AND YIELD PERFORMANCE OF CARROT (*DAUCUS CAROTA* L.) INFLUENCED BY TREATMENTS WITH MICROALGAE EXTRACTS

Irina Sivicka<sup>1\*</sup>, Kaspars Kampuss<sup>1</sup>, Pāvels Semjonovs<sup>2</sup>

<sup>1</sup> Institute of Soil and Plant Sciences, Latvia University of Life Sciences and Technologies

<sup>2</sup> Laboratory of Industrial Microbiology and Food Biotechnology, Institute of Biology, FMLS, University of Latvia

\* Corresponding author. Email: Irina.Sivicka@lbtu.lv

---

The research aimed to the evaluation of growth and yield performance of carrot (*Daucus carota* L.) influenced by treatments with microalgae extracts. At the end of May 2024, carrots (cultivar ‘Nantes 2’ F1, Vilmorin, vegetation period 110 days after germination) were sown in field conditions of the laboratory of Horticulture and Beekeeping of the Latvia University of

Life Sciences and Technologies. From the stage of first leaves (start of June), once per week, total five times plants were sprayed weekly with the solution of ethanol extractions of different microalgae species: *Spirulina*, *Chlorella*, *Tetrademus*, and *Graesiella*. Two concentrations of the extracts were compared with sprays with corresponding ethanol solution as a control. During experiment, no negative effect to carrots' growth and development was noticed for any microalgae extracts. The influence both the type and concentration of microalgae extracts on the growth of carrots was not significant ( $p > 0.05$ ). For yield, kg m<sup>-2</sup>, average result from 3.23 (control variant with drinking water) to 12.03 (*Chlorella*, 10%) was observed. In the case of yield, the significant influence ( $p < 0.05$ ) of concentration was observed, but the influence of the type of microalgae extract was not significant.

**Keywords:** carrot, microalgae, extracts, yield

**Acknowledgements:** this study was performed within the framework of the project No. 22-00-A01612-000014 co-financed by European Agricultural Fund for Rural Development (EAFRD) and supported by the Ministry of Agriculture and Rural Support Service of the Republic of Latvia.

---

## O9: UTILIZING CHEESE WHEY FOR MIXOTROPHIC CULTIVATION OF MICROALGAE *GRAESIELLA EMERSONII*: IMPROVING LIPID AND OMEGA-3 POLYUNSATURATED FATTY ACID CONTENT IN POULTRY DIETS

Kristaps Neiberts\*, Oto Jēkabs Apse, Pāvels Semjonovs

Laboratory of Industrial Microbiology and Food Biotechnology, Institute of Biology, FMLS, University of Latvia, Rīga, Latvija

\* Corresponding author. Email: kristaps.neiberts@lu.lv

---

Photoautotrophic cultivation is the most common method for large-scale microalgal growth. However, this process has some limitations, including long cultivation and low cell productivity due to self-shading at the end of growth (J. Wang et al., 2014). Hence, mixotrophic cultivation method overcomes these drawbacks by combining photosynthesis and organic carbon uptake offering shorter cultivation periods with higher growth rates. This approach minimises self-shading effects and photo limitation by utilising both light and organic compounds as energy source. For instance, dairy wastewater has been utilised as both an energy and a carbon source under mixotrophic conditions to support the growth of various microalgae species, such as *Chlorella vulgaris*, *Chlorella protothecoides*, *Haematococcus pluvialis*, *Scenedesmus obliquus*, *Dunaliella salina* and *Arthrospira platensis* (De Andrade et al., 2022).

Lactose is the main component of cheese whey (CW), contributing to a high chemical oxygen demand (COD) 80–40 g L<sup>-1</sup> and biological oxygen demand (BOD) 30–50 g L<sup>-1</sup>. If not properly managed, CW's high organic content can cause environmental pollution. However, its carbon, nitrogen, and phosphorus content make it a potential substrate for microalgal growth. Effective utilisation requires microalgae capable of metabolising lactose as an organic carbon source (De Andrade et al., 2022). As microalgae are rich in pigments, lipids, polyunsaturated fatty acids, polysaccharides, vitamins, and other compounds, it is essential to determine the cost-effective cultivation method (Ibrahim et al., 2023).

Mixotrophic cultivation combines autotrophic and heterotrophic cultivation methods and has a great option to cut cultivation costs. It has been documented that, in contrast to the cultivation under autotrophic conditions, the cultivation of microalgae under mixotrophic conditions generally improved growth rates, biomass, and other high-value products (Youssef et al., 2024). For instance, in our study, microalgae *Graesiella emersonii* KM01 grown mixotrophically on cheese whey reached  $184.16 \pm 5.75$  mg/g of lipids compared with autotrophic growth, whereas it was estimated only to  $102.10 \pm 1.27$  mg/g.

Microalgal biomass and metabolites are high-value products that serve as alternative feed supplements for broiler chickens due to their rich lipid and Omega-3 fatty acid content. Feeding microalgal biomass can improve body weight and enrich eggs with Omega-3 fatty acids (Abdel-Wareth et al., 2024).

**Keywords:** microalgae, *Graesiella emersonii*, lipids, biomass, poultry diet

**Acknowledgements:** this study was co-financed by the European Agricultural Fund for Rural Development (EAFRD) and supported by the Ministry of Agriculture and Rural Support Service of the Republic of Latvia, grant No. 22-00-A01612-000015.

#### References

1. Abdel-Wareth AaA, Williams AN, Salahuddin M, Gadekar S, Lohakare J. Algae as an alternative source of protein in poultry diets for sustainable production and disease resistance: present status and future considerations. *Front Vet Sci.* 2024;11:1382163.
2. Youssef AM, Gomaa M, Mohamed AKSH, El-Shanawany AA. Enhancement of biomass productivity and biochemical composition of alkaliphilic microalgae by mixotrophic cultivation using cheese whey for biofuel production. *Environ Sci Pollut Res Int.* 2024;31:42875–88.
3. De Andrade AF, Da Costa e Silva PE, De MeloRG, do Nascimento Ferreira MP, Porto ALF, Bezerra RP. Microalgal production under mixotrophic conditions using cheese whey as substrate. *Biol Sci.* 2022;44:8.
4. Wang J, Yang H, Wang F. Mixotrophic cultivation of microalgae for biodiesel production: status and prospects. *Appl Biochem Biotechnol.* 2014;172:3307–29.
5. Ibrahim TNBT, Feisal NAS, Kamaludin NH, et al. Biological active metabolites from microalgae for healthcare and pharmaceutical industries: A comprehensive review. *Bioresour Technol.* 2023;372:128661.

---

## O10: IMPACT OF PYRETHROID INSECTICIDES ON NON-TARGET COLEOPTERA

Zoya Tsybouskaya\*, Austėja Oržekauskaitė

Vytautas Magnus University, Kaunas, Lithuania

\* Corresponding author. Email: zoya.tsybouskaya@gmail.com

---

Insecticides are vital tools for managing pest populations that threaten agricultural productivity and public health by effectively reducing the number of harmful arthropod species. However, their use comes at significant ecological costs (Barathi et al., 2024). In agricultural ecosystems (agrocenoses) and adjacent natural habitats, insecticides disrupt the delicate biological balance, affecting a broad range of non-target organisms. Pyrethroids, widely used insecticides for their effectiveness and relatively low toxicity to mammals, still pose significant

risks to non-target invertebrates (Pathak et al., 2022). The aim of the study was to explore the impact that pyrethroid insecticides have on non-target Coleoptera by using cypermethrin as an example.

We estimated the median lethal dose ( $LD_{50}$ ) of cypermethrin (used as an example compound for pyrethroid insecticides) for each species of Coleoptera studied to understand their sensitivity to pyrethroid insecticides. Furthermore, we investigated the relationship between body size (body mass and body length) of the analysed species of beetles and their sensitivity to pyrethroid insecticides (also using cypermethrin as a model compound).

Laboratory experiments were performed on 24 species of Coleoptera from six different families (Carabidae, Staphylinidae, Silphidae, Histeridae, Aphodiidae, Cetoniidae). The material was collected from forest ecosystems in the vicinity of Kaunas, Lithuania, using standard entomological methods, including: Barber pit-fall traps, manual collection from the litter and soil, butterfly nets. To determine it, the invertebrates (4–8 in one) were put in small closable  $10 \times 6 \times 4$  cm containers (with micro holes in the lids), made of edible plastic. Onto the bottom of the containers, we put cotton disks to absorb surplus moisture, to prevent death by drowning. Using plastic sprayers, one dose (0.37 ml) of an aqueous solution of cypermethrin of a certain concentration was applied (eight different ones were used). The experiment lasted 24 hours. Then the living and dead imagoes of beetles were counted, and  $LD_{50}$  was determined based on data collected.

As a result of the study, we calculated the average lethal doses of cypermethrin for each studied beetle in order to identify the most sensitive species. In addition, we identified a relationship between the body size (weight and body length) of the analysed beetle species and their sensitivity to this insecticide.

Among the analysed beetle groups, Carabidae were found to be the most resistant to cypermethrin. However, exceptions were noted, such as species of the genus *Carabus*, which were very sensitive to cypermethrin. The most sensitive group was Staphylinidae, particularly the species of the genus *Philonthus*. This trend was noted for other groups of dung beetles as well. Cattle dung dwellers were more resistant to cypermethrin than forest litter inhabitants with similar body sizes. It was determined that the taxonomic affiliation of insects is related to the average lethal dose of insecticide. It was also established that for most species of studied beetles there was a relationship between body size and their sensitivity to the insecticide: the shorter the body length and smaller the weight, the higher their sensitivity. However, there are exceptions: coprobionts of the families Histeridae and Aphodiidae were found to be quite resistant to cypermethrin despite their small body size.

**Keywords:** insecticides, pyrethroids, cypermethrin, coleoptera, non-target organisms

#### References

1. Pathak VM, Verma VK, Rawat BS, et al. Current status of pesticide effects on environment, human health and its eco-friendly management as bioremediation: a comprehensive review. *Front Microbiol.* 2022;13:962619.
2. Barathi S, Sabapathi N, Kandasamy S, Lee J. Present status of insecticide impacts and eco-friendly approaches for remediation-a review. *Environ Res.* 2024;240:117432.

---

## O11: THE BURTNIEKU LAKE FLOODPLAIN MEADOWS NATURE RESERVE

Elmīra Boikova\*, Lelde Eņģele, Oskars Keišs, Uvis Suško

*Institute of Biology, FMLS, University of Latvia, Riga, Latvia*

*\* Corresponding author. Email: elmira.boikova@lu.lv*

---

The Latvian Nature Foundation (LDF) has designated wetlands as the habitat of 2024. A wetland is an area that is too wet or covered with a shallow layer of water. Wetlands include floodplain meadows, grass marshes, high marshes, and coastal grasslands. To understand the essence of a wetland, it is necessary to remember that life originated in water, and a wetland is a place where land and water meet and alternate. Therefore, a wetland is full of life. It has the conditions to be able to develop both terrestrial and aquatic species, as well as species that need directly flooded areas. The Burtnieku lake floodplain meadows as a nature reserve and Natura 2000 site covering the area of 432.0 ha was established in 2004. The reserve stretches in a narrow strip around the Burtnieku coast within the territory of the Burtnieku parish. The new nature protection plan for the next 12 years (2025–2037) is in the final stage of development. According to the Habitat Directive, in this territory there are nine biotopes of EU significance, the largest one – Northern boreal alluvial meadows (the type is affected by flooding in spring) – covers 138.0 ha. Compared to the previous nature protection plan for the nesting of 15 specially protected bird species, the population assessment for three species has not changed, for five species the population assessment is currently higher, for three species, it is lower in 2024 than before, – but four species no longer nest in the nature reserve at all: *Gallinago media*, *Tringa stagnatilis*, *Larus minutus*, and *Asio flammeus*. *Crex crex* and *G. media* are of special value for this reserve. Currently, there are only a few suitable places for *C. crex* in the territory of the reserve. The main negative impact on the entire Burtnieku lake meadow reserve is the overgrowth of meadows with bushes, and in some places also with reeds. As a potential enlargement of this protected area with 178.0 ha, Silzemnieku meadows with good conditions for *C. crex* population development is under discussion. The assessment of this territory also illustrates that *Cnidium dubium* can be considered as the second largest and richest place of population distribution after Lubana lake floodplain meadows.

**Keywords:** Natura 2000 site, nature protection plan, population assessment

---



## O12: COMMON STARLING (*STURNUS VULGARIS*) AS A SPECIES FOR CITIZEN SCIENCE AND FUNDAMENTAL RESEARCH

Oskars KeiĶs\*

*Institute of Biology, FMLS, University of Latvia, Riga, Latvia*

*\* Corresponding author. Email: oskars.keiss@lu.lv*

---

The Common Starling (*Sturnus vulgaris*) is a species that the general public sees in its everyday life. In Europe, phenological observations of the arrival of the starling in spring date back to the 19th century. In Latvia, setting up nest boxes for starlings has been popular since 20th century. This makes the common starling almost ideal species for citizen-science projects, involving individuals and schools. The Common Starling in Latvia has been one of the top species ringed between 1925 and 1940 – about 20,000 ringed, and majority of them were ringed by schoolteachers. Until 2024, more than 80,000 common starlings have been ringed in Latvia, and about 1000 recoveries are yielded. During public data gathering on spring arrival dates, the Common Starling has been the most frequent species to be reported due to its synanthropic lifestyle and recognition by the general public. The Common Starling has also been selected as one of the nest-box monitoring species by the Latvian Ornithological Society in the 1980s (Čauns, 1987). It is also selected as Bird of the Year in 2025 by the Latvian Ornithological Society to raise public awareness of the citizen-science contribution to fundamental science project carried out by the Institute of Biology, University of Latvia.

**Keywords:** phenological observations, ringing, ringing recoveries

---

## Poster presentations

## P1: APPLICATION OF MOLECULAR BIOLOGY METHODS FOR THE DETECTION AND GENOTYPING OF *BABESIA CANIS*

Laura Nareckaitė<sup>1</sup>, Miglė Razgūnaitė<sup>1\*</sup>, Justina Snegiriovaitė<sup>1</sup>, Indrė Lipatova<sup>1</sup>,  
Karolina Jankauskaitė<sup>2</sup>, Birutė Karvelienė<sup>2</sup>, Algimantas Paulauskas<sup>1</sup>, Jana Radzijevskaja<sup>1</sup>

<sup>1</sup> Vytautas Magnus University, Kaunas, Lithuania

<sup>2</sup> Lithuanian University of Health Sciences, Kaunas, Lithuania

\* Corresponding author. Email: migle.razgunaite@vdu.lt

Canine babesiosis is a widespread tick-borne infectious disease caused by intraerythrocytic protozoa *Babesia canis*. Over the past two decades, the number of cases of canine babesiosis in Europe has been increasing significantly, and higher numbers of cases have been observed in Lithuania as well. As the disease spreads rapidly, greater attention is needed for accurate diagnosis and effective treatment. Diagnostic methods commonly used in veterinary clinics, such as blood smears, are not always sensitive enough to identify the infection or do not allow for accurate identification of the pathogen. We aimed to investigate the genetic diversity of *B. canis* strains isolated from infected dogs in Lithuania using sequence analysis of *B. canis* 18S rRNA and Bc28.1 gene fragments. Blood samples were collected from 107 symptomatic dogs in veterinary clinics from spring to autumn of 2024. PCR tests were performed for both 18S rRNA and Bc28.1 genes. *B. canis* DNA was detected in 62 samples (66.34%) out of 107. Sequence analysis of the 18S rRNA gene fragment showed two variable sites in 81–82 nucleotide positions, dividing *B. canis* into three different genotypes. Bc28.1 gene fragment analysis identified four gene sequence variants, which were classified into three genotypes based on two nucleotide polymorphisms. The results highlight discrepancies between diagnoses made in veterinary clinics using blood smears and those obtained through molecular methods. Furthermore, they demonstrate the potential of molecular techniques for detecting genetic variability among *B. canis* strains, which could improve diagnostic accuracy and epidemiological surveillance.

**Keywords:** canine babesiosis, *Babesia canis*, 18S rRNA gene, Bc28.1 gene

**Acknowledgements:** this study was supported by the Research Council of Lithuania (Grant No. S-MIP-23/19).

---

## P2: MOLECULAR IDENTIFICATION OF *BORRELIA BURGDORFERI* SENSU LATO IN TICKS IN KAUNAS CITY, LITHUANIA

Saulius Bernotas\*, Justina Snegiriovaite, Migle Razgūnaitė, Indrė Lipatova, Algimantas Paulauskas, Jana Radzijeuskaja

Research Institute of Natural and Technological Sciences, Vytautas Magnus University, Kaunas, Lithuania

\* Corresponding author. Email: saulius.bernotas@vdu.lt

---

Lyme disease, caused by *Borrelia burgdorferi* sensu lato (s. l.), is one of the most prevalent tick-borne infections in Europe. While the disease is well-documented, data on *Borrelia* prevalence in urban and peri-urban environments, including Lithuania, remains limited. Since ticks serve as key vectors, monitoring their infection rates is essential for assessing the risk of Lyme disease in these settings. The study aimed to identify *B. burgdorferi* (s. l.) in ticks collected in Kaunas city, Lithuania, using molecular methods. A total of 584 questing *Ixodes ricinus* ( $n = 527$ ) and *Dermacentor reticulatus* ( $n = 57$ ) ticks were collected by flagging in various urban and peri-urban areas of Kaunas city in 2023–2024. The presence of *Borrelia* DNA in ticks was screened using real-time PCR targeting the 23S rRNA gene. For identification of *B. burgdorferi* s. l. species, the partial outer surface protein A (ospA) gene, the chromosomal flagellin gene, and the 16S (rrs)–23S (rrlA) rRNA intergenic spacer (IGS) region were amplified using conventional and nested PCR assays, and sequenced. Overall, *Borrelia* DNA was detected in 20.72% (121/584) of the collected ticks. Infection was found only in *I. ricinus*. Three *Borrelia* species belonging to the *B. burgdorferi* s. l. complex were identified: *B. burgdorferi sensu stricto*, *B. garinii*, and *B. afzelii*. Our results demonstrate the potentially high human risk of exposure to tick-borne infection with *B. burgdorferi* s. l. in cities in Lithuania.

**Keywords:** *Borrelia burgdorferi* s. l., ticks, urban, peri-urban, Lithuania

**Acknowledgements:** this study was supported by the Research Council of Lithuania (Grant No. S-MIP-23/19).

### References

1. Trevisan G, Cinco M, Trevisini S, di Meo N, Chersi K, Ruscio M, Forcione P, Bonin S. *Borreliae* part 1: *Borrelia* Lyme group and *Echidna*-reptile group. *Biology*. 2021;10:1036.
  2. Marques AR, Strle F, Wormser GP. Comparison of Lyme disease in the United States and Europe. *Emerg Infect Dis*. 2021;27:2017.
  3. Hornok S, Meli ML, Gönczi E, Halász E, Takács N, Farkas R, Hofmann-Lehmann R. Occurrence of ticks and prevalence of *Anaplasma phagocytophilum* and *Borrelia burgdorferi* s.l. in three types of urban biotopes: forests, parks and cemeteries. *Ticks Tick Borne Dis*. 2014;5:785–9.
  4. Snegiriovaite J, Radzijeuskaja J, Lipatova I, Razgūnaitė M, Paulauskas A. The distribution of Ixodid ticks and their infection with *Borrelia* spp. in urban and periurban parks in Lithuania. *Parasites of 21 Century XV. Slovak and Czech Parasitological Days, Tesáre – Nové Mlýny, Slovakia, 13–17 May 2024: abstract book*.
-

## P3: MOLECULAR SCREENING FOR TICK-BORNE ENCEPHALITIS VIRUS IN TICKS FROM URBAN ENVIRONMENTS

Martynas Jonkus<sup>1</sup>, Miglė Razgūnaitė<sup>1\*</sup>, Justina Snegiriovaitė<sup>1</sup>, Indrė Lipatova<sup>1</sup>,  
Arūnas Stankevičius<sup>2</sup>, Algimantas Paulauskas<sup>1</sup>, Jana Radzijeuskaja<sup>1</sup>

<sup>1</sup> Vytautas Magnus University, Faculty of Natural Sciences, Kaunas, Lithuania

<sup>2</sup> Lithuanian University of Health Sciences, Kaunas, Lithuania

\* Corresponding author. Email: migle.razgunaite@vdu.lt

Climate change has significantly influenced tick ecology by extending their seasonal activity and facilitating their expansion into new habitats. As a result, the incidence of tick-borne diseases is rising in previously low-risk areas, posing increasing public health concerns. Among these, tick-borne encephalitis (TBE) is one of the most significant vector-borne diseases in Europe, with the Baltic countries accounting for nearly one-third of reported European cases. This study assessed the prevalence of tick-borne encephalitis virus (TBEV) in questing ticks in Lithuania and key risk factors for transmission. Between 2023 and 2024, 544 ticks (*Ixodes ricinus*,  $n = 525$ ; *Dermacentor reticulatus*,  $n = 19$ ) were collected using the flagging method in urban areas and grouped into 47 pools based on species and developmental stages. TBEV detection was performed using real-time RT-PCR and virus isolation in two cell lines (Neuro-2a, MARC-145). TBEV RNA was detected in 15.26% (83/544) of samples. Virus isolation proved more effective than direct RT-PCR, with a total minimum infection rate (MIR) of 4.41% across both cell lines, compared to 1.47% from direct RT-PCR. The MIR for *I. ricinus* was 1.3% (7/525) and for *D. reticulatus* 5.3% (1/19). Positive samples included six nymphs, nine females, and seven males, highlighting widespread TBEV circulation across developmental stages and sexes. These findings suggest that *D. reticulatus* may play a greater role in TBEV transmission than previously recognised. The study underscores the need for ongoing tick surveillance and public health strategies, as well as the importance of virus isolation for improving detection accuracy.

**Keywords:** tick-borne encephalitis, *Ixodes ricinus*, *Dermacentor reticulatus*, questing ticks

**Acknowledgements:** this study was supported by the Research Council of Lithuania (Grant No. S-MIP-23/19).

### References

1. Sidorenko M, Radzijeuskaja J, Mickevičius S, Bratchikov M, Mardosaitė-Busaitienė D, Sakalauskas P, Paulauskas A. Phylogenetic characterisation of tick-borne encephalitis virus from Lithuania. PLoS One. 2024;19:e0296472.
2. Sidorenko M, Radzijeuskaja J, Mickevičius S, Bratčikovienė N, Paulauskas A. Prevalence of tick-borne encephalitis virus in questing *Dermacentor reticulatus* and *Ixodes ricinus* ticks in Lithuania. Ticks Tick Borne Dis. 2021;12:101594.
3. Pautienius A, Dudas G, Simkute E, Grigas J, Zakiene I, Paulauskas A, Armonaite A, Zienius D, Slyzius E, Stankevicius A. Bulk milk tank samples are suitable to assess circulation of tick-borne encephalitis virus in high endemic areas. Viruses. 2021;13:1772.

---

## P4: DEVELOPMENT OF COST-EFFECTIVE MOLECULAR METHOD TO DISCRIMINATE DIFFERENT *LISTERIA MONOCYTOGENES* STRAINS IN FOOD PROCESSING SECTOR

**Donata Grikšaitė\*, Dalius Butkauskas**

*Nature Research Centre, Vilnius, Lithuania*

*\* Corresponding author. Email: grikstaitedonata@gmail.com*

---

*Listeria monocytogenes* is a common environmental bacterium associated with natural and artificial-human made environments such as soil, water, vegetation, animals, and food processing environments. This microorganism is characterised by high genetic diversity, which determines its ability to adapt to various environmental conditions and cause disease in various animal and human populations. Knowledge of the genetic diversity of strains circulating in food processing sectors is essential to assess the risks associated with this pathogen and improve food safety, therefore the genetic diversity of *L. monocytogenes* in the environment plays a very important role in understanding its transmission routes. This study investigates the genetic diversity of *L. monocytogenes* strains in food processing environments. During the study, 27 samples were collected from two food processing plants after environmental specimens were propagated in ALOA and OXFORD media, and their DNA was subsequently isolated. Using the PCR (Polymerase Chain Reaction) method, *L. monocytogenes* was detected in 27 cases. Using an advanced sequencing method – multi-locus sequence typing (MLST), we determined the genetic profiles of the isolates. This methodology allows typing of bacteria by several specific gene loci, providing the possibility to identify different bacterial strains. Our analysis revealed the diversity of isolates from the production environment, i.e., three sequence types representing three bacterial strains were identified. It is important to note that two types were repeated in both food processing plants, while one type was exclusively identified in one food processing environment.

**Keywords:** DNA studies, genetic diversity, listeriosis, PCR

### References

1. Orsi RH, den Bakker H., Wiedmann M. *Listeria monocytogenes* lineages: genomics, evolution, ecology, and phenotypic characteristics. *Int J Med Microbiol.* 2011;301:79–96.
  2. Liu D. *Handbook of Listeria monocytogenes*. CRC Press: 2008. 512 p.
-

---

## P5: THE ROLE OF ANTHROPOGENIC PRESSURES IN SHAPING THE GENETICS OF RED DEER (*CERVUS ELAPHUS*)

Evelina Genevičienė, Irma Ražanskė, Loreta Gričiuvienė, Kastytis Šimkevičius, Artūras Kibiša, Algimantas Paulauskas\*

Vytautas Magnus University, Kaunas, Lithuania

\* Corresponding author. Email: algimantas.paulauskas@vdu.lt

---

Native to Central Europe, red deer have expanded their range over the past thirty years and are now inhabiting areas where they coexist with the introduced sika deer. This overlap has led to several ecological challenges, particularly hybridisation between the two species and changes to their genetic structure. During the rut, aggressive sika males may attack red deer stags and mate with red deer hinds, raising the chances of hybridisation and gene flow between the species. Research into the genetic diversity of both red and sika deer was conducted in Lithuania using tissue samples from legally hunted animals. Genotyping was performed based on nuclear DNA microsatellite loci (STR), and when combined with statistical analysis, the data provided a clearer understanding of the populations and their genetic differences. Hybridisation cases between the two species have been documented in several EU countries, highlighting the growing interaction between red and sika deer. Additionally, the study examined the impact of sika deer on red deer populations, particularly due to human activities, such as hunting and habitat disruption. This research emphasizes the importance of monitoring and managing deer populations to understand the broader ecological implications of these interspecies interactions. The increasing presence of sika deer in regions traditionally inhabited by red deer poses significant challenges for conservation efforts. As hybridisation becomes more common, it raises concerns about the potential loss of unique genetic traits in red deer populations

**Keywords:** Red deer, genetic diversity, anthropogenic impact

### References

1. Biedrzycka A, Solarz W, Okarma H. Hybridization between native and introduced species of deer in Eastern Europe. *J Mammalogy*. 2012;93:1331–41.
  2. Ražanskė I, Gibiežaitė JM, Paulauskas A. Genetic analysis of red deer (*Cervus elaphus*) and sika deer (*Cervus nippon*) to evaluate possible hybridisation in Lithuania. *Baltic Forestry*. 2017; 23: 683–90.
-

---

## P6: GENETIC DIVERSITY OF *BARTONELLA* SPP. IN LITHUANIAN BATS BASED ON *GLTA* GENE SEQUENCES

Paulina Maceinaitė\*, Povilas Sakalauskas, Algimantas Paulauskas

Vytautas Magnus University, Kaunas, Lithuania

\* Corresponding author. Email: paulina.maceinaite@stud.vdu.lt

---

Analysing the genetic diversity of *Bartonella* in bats is essential in assessing the risks associated with its transmission. In addition, our understanding of host specificity and potential pathogenicity is improving with the discovery of new genetic lineages of *Bartonella*. As some *Bartonella* species can cause human diseases such as bartonellosis, this study is important for epidemiological surveillance and prevention of zoonotic diseases. Bats are one of the most diverse and widespread mammalian species. They are important for ecosystems. They can be a factor in disease transmission due to their ability to adapt to different environments and feeding habits. To study bats as *Bartonella* reservoirs, samples collected in Lithuania from 2018 to 2020 were analysed. DNA was extracted from lung, heart, kidney, and spleen tissues using the Genomic DNA Purification Kit. *Bartonella* spp. detection was performed using real-time PCR targeting the *gltA* gene, followed by agarose gel electrophoresis, purification, and sequencing. Sequences were compared against the GenBank database, and phylogenetic relationships were determined by maximum likelihood using the MEGA 11 software. A total of 53 bats from five species (*Myotis daubentonii*, *Eptesicus serotinus*, *Pipistrellus nathusii*, *Vespertilio murinus*, and *Nyctalus noctula*) were analysed. Five *Bartonella* samples were subjected to sequencing, yielding sequences with 49 variable nucleotide positions. Comparative analysis with GenBank reference sequences revealed a similarity range of 94% to 100%. The sequences were grouped into two clusters by phylogenetic analysis: one was most closely related to *Bartonella phoceensis*, which was previously detected in rodents from Malaysia, and *Bartonella alsatica*, which was previously found in *Apodemus speciosus* rodents, and the other was related to *Bartonella bacilliformis*, which was found in a human from Peru. Phylogenetic analysis showed that there are various species of *Bartonella* in Lithuanian bats, which highlights the role of bats as natural reservoirs.

**Keywords:** bats, *Bartonella*, *gltA* gene, host specificity

### References

1. Banerjee A, Baker ML, Kulcsar K, Misra V, Plowright R, Mossman K. Novel insights into immune systems of bats. *Front Immunol.* 2020;11:26.
  2. Urushadze L, Bai Y, Osikowicz L, et al. Prevalence, diversity, and host associations of *Bartonella* strains in bats from Georgia (Caucasus). *PLoS Negl Trop Dis.* 2017;11(4):e0005428.
  3. Vayssier-Taussat M, Moutailler S, Féménia F, et al. Identification of novel zoonotic activity of *Bartonella* spp., France. *Emerg Infect Dis.* 2016;22(3):457–62.
-

---

## P7: APPLICATION OF ENZYME-LINKED IMMUNOSORBENT ASSAY FOR THE DETECTION OF ANTIBODIES AGAINST THE MEASLES VIRUS

Justina Dagytė\*, Indrė Lipatova, Algimantas Paulauskas

*Vytautas Magnus University, Kaunas, Lithuania*

\* Corresponding author. Email: justina.dagyte@vdu.lt

---

Measles is a highly contagious, acute, and severe disease that affects people of all ages (Moss, 2017). It causes fever, maculopapular rash and upper respiratory tract inflammation (Fisher et al., 2014). The infection is extremely dangerous for older adults with a weak immune system and children (Bester, 2016; Moss, 2017). Despite universal vaccination programmes, the outbreaks of measles have been increasing in recent years, suggesting gaps in immunity, potentially due to waning immune memory or insufficient vaccine coverage (Patel et al., 2020). This study aimed to evaluate the immune status of adults and to identify individuals at risk of measles infection in Lithuania. The enzyme-linked immunosorbent assay method applied in this research allowed to detect levels of IgG antibodies in the participants. Serum samples from a total of 88 individuals were included in this study. The results revealed that the majority of individuals with insufficient IgG antibody levels were from the 1976–1981 and 1982–1987 age groups. Participants with negative results were either uncertain about their vaccination status or had received only a single vaccine dose. The distribution of these results was similar between sexes. These findings highlight the importance of improving vaccination coverage, particularly for individuals with incomplete immunisation histories, in order to reduce the risk of the outbreaks of measles.

**Keywords:** measles, IgG antibodies, ELISA, Lithuania

### References

1. Bester JC. Measles and measles vaccination. *JAMA Pediatrics*. 2016;170:1209–15.
  2. Fisher DL, Defres S, Solomon T. Measles-induced encephalitis. *QJM*. 2015;108:177–82.
  3. Moss WJ. Measles. *Lancet*. 2017;390:2490–502.
  4. Patel MK, Antoni S, Nedelec Y, Sodha S, Menning L, Ogbuanu IU, Gacic Dobo M. The changing global epidemiology of measles, 2013–2018. *J Infect Dis*. 2020;222:1117–28.
- 

## P8: APPLICATION OF MOLECULAR METHODS FOR THE DETECTION OF LYME DISEASE AGENTS

Ugnė Medikaitė\*, Indrė Lipatova, Justina Snegiriovaitė, Jana Radzijeuskaja,  
Algimantas Paulauskas

*Vytautas Magnus University, Lithuania*

\* Corresponding author. Email: ugne.medikaite@vdu.lt

---

Lyme disease is the most common vector-borne disease in North America and Europe. This infection is caused by the spirochete bacteria complex *Borrelia burgdorferi* sensu lato and is transmitted to humans from the bite of an infected tick (Marques et al., 2021; Radolf et al., 2021).



The increased incidence of Lyme disease over recent decades could be linked to factors such as climate and environmental changes, the growing distribution of ticks and their hosts (Septfons et al., 2021). Understanding the role of potential reservoir hosts in disease transmission is required for effective public health strategies. Red squirrels (*Sciurus vulgaris*), which are commonly found in natural, peri-urban, and urban areas where humans frequently interact with wildlife, may act as a reservoir host for maintaining *Borrelia* spp. (Krauze-Gryz et al., 2015). In this study, molecular methods were applied for the detection of *Borrelia* DNA in red squirrels from urban parks in Kaunas city, Lithuania. A total of 45 red squirrel tissue samples were analysed using real-time PCR with *Borrelia*-specific primers. Positive samples were further analysed through PCR amplification targeting the intergenic spacer (IGS) region, *ospA*, and *fla* genes, followed by sequencing. The results revealed that 37.7% of the tested samples were positives for *Borrelia* DNA. The detection rates varied among genes, with *ospA* detected in 17.7% of samples, IGS in 20.0%, and *fla* in 37.7%. Three pathogenic species were identified by sequencing: *B. afzelii*, *B. garinii*, and *B. burgdorferi* sensu stricto. These findings indicate that red squirrels could play a role in the spread of Lyme disease by hosting multiple *Borrelia* species. This highlights the need for ongoing monitoring of wildlife and tick populations in areas where the disease is common.

**Keywords:** *Borrelia burgdorferi* sensu lato, spirochete bacteria complex, disease transmission

#### References

1. Radolf JD, Strle K, Lemieux JE, Strle F. Lyme Disease in Humans. *Curr Issues Mol Biol.* 2021;42:333–84.
2. Septfons A, Figoni J, Gautier A, Soullier N, de Valk H, Desenclos JC. Increased awareness and knowledge of Lyme Borreliosis and tick bite prevention among the general population in France: 2016 and 2019 health barometer survey. *BMC Public Health.* 2021;21:1808.
3. Marques AR, Strle F, Wormser GP. Comparison of Lyme Disease in the United States and Europe. *Emerg Infect Dis.* 2021; 27(8):2017–24.
4. Krauze-Gryz, D. A. G. N. Y., Gryz, J. A. K. U. B. A review of the diet of the red squirrel (*Sciurus vulgaris*) in different types of habitats. In: Red squirrels: ecology, conservation & management in Europe. European Squirrel Initiative: Stoneleigh Park, Warwickshire CV8 2LG, UK; 2015. p. 39–50.

---

## P9: PRELIMINARY RESULTS ON MUSCLE TISSUES PARASITES OF MUSTELIDS IN LITHUANIA

**Ramunė Tamošiūnaitė\*, Evelina Maziliauskaitė, Petras Prakas, Dalius Butkauskas**

*Nature Research Centre, Vilnius, Lithuania*

\* Corresponding author. Email: ramune.tamosiunaite@gmc.stud.vu.lt

---

Mustelids are found worldwide and occupy various habitats, including forests, grasslands, wetlands, and urban areas. Their wide variety of food often exposes them to infected wildlife, which makes them reservoirs of zoonotic infections. Mustelids frequently transmit various nematodes, for instance, *Trichinella* spp. To ensure the safety of food it is necessary to investigate these mammals because they can be potential vectors of *Trichinella* spp. parasites. Furthermore, mustelids carry a variety of trematodes species such as *Alaria* spp., *Neodiplostomum* spp., and *Strigea* spp. Therefore, the objectives of the present study were to establish parasite infection prevalence and to use molecular methods to identify parasite species. Sixty mammals of the Mustelidae family – 26 European pine marten, 19 American mink, 13 European polecats, one European badger,

and one stone marten – were collected from different regions of Lithuania. The active artificial method was used to investigate the infection prevalence and intensity of *Trichinella* spp. in muscle samples. The microscopic compressor method was used to detect different trematodes metacercariae. *Trichinella* species were identified using multiplex-PCR, while trematodes were identified by ITS2 region and Sanger sequencing of amplified fragments. Seven of the 60 mustelids were positive for at least one parasite. Prevalence of *Trichinella* spp. and trematodes were 6.67% and 5.00%, respectively. Overall, 40 larvae were isolated for molecular confirmation of *Trichinella* species. The 30 larvae isolated from one European polecat and two European pine martens were identified as *T. britovi* being the dominant *Trichinella* species in wild animals in Lithuania. Also, we found mixed *T. britovi*/*T. spiralis* in one positive American mink (ratio 8:2). One of the three trematode-infected animals was examined by molecular analysis, which showed 100% identity to *Neodiplostomum spathula*. In addition, for the first time *Trichinella* spp. was found in American mink and *Neodiplostomum spathula* was detected in Lithuania.

**Keywords:** mustelids, *Trichinella* spp., *Neodiplostomum* spp., molecular identification

---

## P10: SARCOCYSTIS SPECIES RICHNESS IN THE DIAPHRAGM AND MYOCARDIUM OF ALPINE IBEX (*CAPRA IBEX*)

Eglė Rudaitytė-Lukošienė\*, Petras Prakas, Dalius Butkauskas

Nature Research Centre, Vilnius, Lithuania

\* Corresponding author email: egle.rudaityte@gmail.com

---

The cyst-forming coccidia of the genus *Sarcocystis* (Apicomplexa: Sarcocystidae) are widespread parasites of mammals, particularly domestic and wild ruminants. These protozoan parasites form sarcocysts in muscle tissues of intermediate hosts. Research on *Sarcocystis* in wild members of the subfamily Caprinae is, however, rather limited. A comprehensive investigation of sarcocysts in Alpine ibex (*Capra ibex*) was previously conducted using solely morphological techniques. In the period of 2021–2024, diaphragm and myocardium muscle samples of eight Alpine ibex from Austria were examined for sarcocysts. The detected sarcocysts were morphologically described using light microscopy and transmission electron microscopy. The cysts were classified into three morphological types (I–III), distinguished by their distinct morphometric parameters, including the thickness of the cyst wall and the presence of finger-like or hair-like protrusions, respectively. Furthermore, molecular identification and characterisation of individual sarcocysts was conducted using the 18S rRNA gene and *cox1* sequencing, followed by conducting phylogenetic analyses on the resulting sequences. This study offers a significant advancement in our understanding of *Sarcocystis* species diversity, as it is the first to employ molecular methods to identify this parasite in Alpine ibex. The analysis revealed that type II sarcocysts exhibited the closest phylogenetic relationship with *Sarcocystis* species transmitted by corvid birds, while types I and III showed a closer relationship with species transmitted by canids. However, the 18S rDNA, in contrast to the *cox1*, was not sufficiently variable to accurately delineate the phylogenetic relationships of numerous *Sarcocystis* species. This study corroborates the role of the Alpine ibex as a host of three *Sarcocystis* species.

**Keywords:** *Sarcocystis*, Alpine ibex, molecular identification, *cox1*, phylogeny

---

---

## P11: GENETIC CHARACTERISATION OF *SARCOCYSTIS* CYSTS ISOLATED FROM MUSCLES AND BRAIN OF VOLES FROM LITHUANIA

Dovilė Laisvūnė Bagdonaitė\*, Eglė Rudaitytė-Lukošienė, Petras Prakas, Dalius Butkauskas

Laboratory of Molecular Ecology, Nature Research Centre, Vilnius, Lithuania

\* Corresponding author. Email: dovile.bagdonaitė@gamtc.lt

---

*Sarcocystis* (Apicomplexa) is a genus of parasites that employs a variety of hosts—mammals, birds, or reptiles to complete their life cycle. These parasites are characterised by the formation of sarcocysts in the muscles or CNS of intermediate hosts and the development of sporocysts in the intestines of definitive hosts. To date, over 40 different *Sarcocystis* species have been identified in rodents. Majority of the investigations have focused on synanthropic rodent species, such as the house mouse (*Mus musculus*) and brown rat (*Rattus norvegicus*), thus data concerning prevalence and diversity of *Sarcocystis* species in wild mice and voles remain sparse. Due to the limited amount of available data, the precise effects of these parasites to their hosts remain to be fully determined. However, it is known that certain *Sarcocystis* species, which utilise rodents as intermediate hosts, are either pathogenic or increase the risk of predation of affected animals. In recent years, molecular techniques have been increasingly used to distinguish even morphologically similar *Sarcocystis* species and to study animals for which previous studies were too difficult due to technical barriers. In this study, we aimed to molecularly characterise *Sarcocystis* parasites isolated from the common vole (*Microtus arvalis*) and the bank vole (*Clethrionomys glareolus*).

Muscle and brain samples of 52 common voles and bank voles collected in different regions of Lithuania were examined for *Sarcocystis* spp. Parasites were detected and isolated from tissues using light microscopy. *Sarcocystis* species were identified and characterised within 18S rRNA, 28S rRNA, ITS1, ITS2, cox1, cytb and rpoB. Three sarcocysts were found in the muscles of a single common vole. These sarcocysts were morphologically similar to those of the recently described *Sarcocystis myodes* species in Lithuania. Molecular analysis of the four gene markers revealed that sarcocysts belong to genetically new *Sarcocystis* species. For the description of a new *Sarcocystis* species, a more detailed morphological analysis of sarcocysts involving transmission electron microscopy is needed. Furthermore, *S. glareoli* cysts found in the brain samples of the bank vole were genetically characterised using seven genetic markers located in nuclear, mitochondrial, and apicoplast genomes. The results obtained in this study are important for the development of molecular identification of parasites of the genus *Sarcocystis* and for the overall understanding of the species diversity of *Sarcocystis* parasites in wild small mammals.

**Keywords:** *Sarcocystis*, voles, brain, muscles, molecular characterisation

---

---

## P12: OPTIMIZATION AND CHALLENGES IN DNA EXTRACTION FROM *TETRAO UROGALLUS* FEATHERS

Evelina Slavnova\*, Dace Grauda, Dalius Butkauskas

*Institute of Biology, FMLS, University of Latvia*

\* Corresponding author. Email: linamilss28@gmail.com

---

The western capercaillie (*Tetrao urogallus*) is a key species in boreal forests, serving as a biodiversity indicator in Latvia and Lithuania (Bagrađe et al., 2017). However, habitat fragmentation, population decline, and reproductive isolation threaten genetic diversity and survival of capercaillie (Vallant et al., 2018). To assess genetic diversity of declining capercaillie populations in Latvia and Lithuania, the non-invasive method based on collection of feathers for DNA extraction as the most suitable option was chosen in attempts not to disturb birds, especially during breeding season. To optimise DNA extraction from collected *T. urogallus* feathers, we used muscle tissue from two dead birds as control specimens and down feathers collected in natural habitat or obtained from capercaillie breeding farm, located in Viešvilē Nature Reserve (Lithuania). Initial extractions from feathers and muscles yielded high DNA concentrations but contained impurities that affected PCR performance. To improve quality, we developed a two-step purification protocol. After initial purification, samples were air-dried and subjected to a 70% ethanol wash, significantly enhancing DNA purity. Agarose gel electrophoresis and PCR analysis confirmed successful primer binding after purification, demonstrating the effectiveness of the method. Further modifications of the DNA extraction and PCR protocols improved amplification efficiency, leading to the collection of amplified D-loop fragments subjected to sequencing in the Laboratory of Molecular Ecology of Nature Research Centre. The non-invasive genetic studies of *T. urogallus* are going to be continued including sequencing of informative mtDNA fragments and analysis of microsatellite loci. Our optimised method provides a reliable approach for obtaining high-quality DNA from feathers, supporting future genetic research and conservation efforts. The optimised protocol can also be applied to other bird species requiring non-invasive sampling. Additionally, integrating this method with sequencing of D-loop region or analysis of some nuclear loci could offer deeper insights into genetic diversity, population structure, and adaptive potential. These advancements will aid conservation strategies, including habitat management and genetic rescue efforts.

**Keywords:** Western Capercaillie, DNA extraction, non-invasive sampling, PCR optimization

**Acknowledgements:** we would like to express our sincere gratitude to Algis Butleris and Kristina Butlerienē from Viešvilē Nature Reserve for providing feather and muscle material, and Aivars Ornicāns, a scientist in Silava, for providing feather material from Latvia.

### References

1. Bagrađe G, Baumanis J, Bitenieks K, Done G, Lūkins M, Ornicāns A, Ozoliņš J, Pilāte D, Priediti A, Stepanova A, Šuba J, Žunna A. Medņu aizsardzībai nozīmīgo vides faktoru izpēte. Silava. 2017;1–73.
  2. Vallant S, Niederstätter H, Berger B, Lentner R, Parson W. Increased DNA typing success for feces and feathers of capercaillie (*Tetrao urogallus*) and black grouse (*Tetrao tetrix*). Ecol Evol. 2018;8:3941–51.
-

## P13: BREEDING OF TETRAPLOID RED CLOVER IN LATVIA

Aija Rebane<sup>1\*</sup>, Dace Grauda<sup>2</sup>, Paula Marta Muceniece<sup>2</sup>, Sarmite Rancane<sup>1</sup>

<sup>1</sup> Institute of Agriculture, Latvia University of Life Sciences and Technologies Agriculture, Skriveri, Latvia

<sup>2</sup> Institute of Biology, FMLS, University of Latvia; Riga, Latvia

\* Corresponding author. Email: aijarebane@inbox.lv

Increasing the sowing area of legumes, including red clover (*Trifolium pratense* L.), is an important task in modern agriculture not only in Latvia, but in many European and other countries. Growing legumes gives the opportunity to maintain and increase soil fertility, obtain cheap and comprehensive feed by decreasing the use of nitrogen fertiliser, and preserve nature. The ability of legumes, including red clover, to fix atmospheric nitrogen is particularly relevant today, given the drastically rising energy prices. Our feed producers are becoming more demanding and interested in new, more productive varieties. Therefore, breeding work should not be interrupted or stopped due to the rapid changes in growing and harvesting technologies, climatic conditions, diseases, and pests. The breeding goal is to offer consumers a variety with high adaptive potential, able to function effectively in different soil types, which can quickly build a large photosynthetic surface. Competitive struggle with weeds for space and nutrients, in the form of large, stable, and biomass and seed yield, winter-hardy, persistent, and resistant to diseases and pests – they are the main requirements for a new variety. Worldwide, red clover breeding has focused mainly on the creation of genotypes with higher biomass yield and increased adaptability. To meet consumer demands in these changing conditions, breeders need to work continuously using both classic and modern breeding methods.

Since the 1960s, red clover breeding programs pay special attention to the development of tetraploid varieties (Jansone et al., 2014). Tetraploid red clover is characterised by better ecological adaptability, higher biomass, and disease resistance compared to diploid plants. Treatment of diploid sprouts with colchicine is commonly used to obtain red clover polyploids, resulting in a change in the number of chromosomes (Slater et al., 2003). However, a relatively small number of stable tetraploids can be obtained using this method. To facilitate the acquisition of tetraploid red clover breeding material, various *in vitro*, molecular, cytometric, and flow cytometric methods were used in collaboration with researchers from the Environmental Genetics Laboratory of the Institute of Biology of the University of Latvia. Seeds of medium-early red clover cultivars 'Dižstende', 'Stendes agrais', and 'Jancis' were used to obtain tetraploid plants.

As part of the red clover breeding programme of the Institute of Agriculture of the Latvian University of Life Sciences and Technologies (LBTU IA), in cooperation with researchers from the Plant Genetics Laboratory of the Institute of Biology of the University of Latvia, 390 samples of red clover plants were submitted for further evaluation between 2013 and 2017. The plants were planted in open field conditions in the Skriveri Institute (LBTU IA) experimental fields and evaluated for six generations. From the selection and family nurseries, only the two most promising samples were collected, which will continue to be used in the future breeding programme. To make sure that the valuable qualities for which the samples were selected are passed on to the future, it is important to re-sow, establish, and evaluate them in progeny test nurseries.

**Keywords:** tetraploid red clover, *in vitro*, molecular methods, flow cytometry

## References

1. Jansone B, Rancāne S, Jansons A, Rebāne A, Jermuša G. Breeding of perennial legumes at the Agricultural Research Institute in Skrīveri. Proceedings of scientific-practical conference 'Līdzsvarota Lauksaimniecība', Jelgava, 2014. p. 81–86. Latvian.
2. Slater A, Nigel W, Scott M, Fowler R. Plant Biotechnology: The Genetic Manipulation of Plants. Oxford: Oxford University Press, 2003, 368 p.

# P14: ASSESSMENT OF ANTIMICROBIAL PROPERTIES OF 3D BIOTEXTILE WITH INCORPORATED NANOPARTICLES

Elina Ažēna<sup>1</sup>, Dace Grauda<sup>1\*</sup>, Valters Gobins<sup>1</sup>, Inga Lasenko<sup>3</sup>, Dalius Butkauskas<sup>2</sup>

<sup>1</sup> Institute of Biology, FMLS, University of Latvia, Riga, Latvia

<sup>2</sup> Nature Research Centre, Vilnius, Lithuania

<sup>3</sup> JLU Technologies Ltd, Riga, Latvia

\* Corresponding author. Email: dace.grauda@lu.lv

Antimicrobial activity was evaluated for fabrics incorporating succinite, silica, aluminium oxide, silver nanoparticles, and fabrics treated with crystalline succinic acid. For the studies, microorganism cultures (University of Latvia, Microbial Strain Collection of Latvia) representing human microbiota were used, including gram-positive and gram-negative bacteria and fungi. *Escherichia coli* (MSCL332) and *Enterococcus faecalis* (MSCL302) served as indicators of faecal contamination, while *Staphylococcus aureus* (MSCL334) and *Candida albicans* (MSCL378) were included as opportunistic pathogens linked to skin infections. Antimicrobial activity was assessed using the disk diffusion and parallel streak methods, which evaluate microbial growth inhibition on agar. Additional tests determined the impact of fabrics on microbial growth in a liquid medium, analysed via optical density, colony-forming unit (CFU) counts, and flow cytometry. A BD FACSJazz® Cell Sorter (BD Biosciences, USA) was used for flow cytometry to measure the relative fluorescence of bacterial cells. A 488 nm blue laser excited cell fluorescence, and the emission was measured at the 585 nm channel. The flow cytometer was calibrated using Sphero™ rainbow calibration particles (3.0 µm, BD Biosciences, USA) in PBS solution before measurements. A coefficient of variation (CV) below 3% was considered a successful calibration. The intensity of bacterial relative fluorescence was expressed in arbitrary logarithmic units. For each sample, events were recorded over a fixed 3-min period.

The disk diffusion method demonstrated antimicrobial activity in succinic acid-treated fabrics, showing inhibition zones for *S. aureus*, *E. faecalis*, and *E. coli*, but not for *C. albicans*. The parallel streak method confirmed these results. Further evaluation showed decrease in bacterial relative fluorescence of *E. coli* and *S. aureus* colonies grown in liquid medium with succinic acid-treated fabrics. Additionally, antimicrobial activity was observed in bacterial colonies grown with fabrics containing succinic nanoparticles.

**Keywords:** biotextile, succinite nanoparticles, succinic acid, antimicrobial activity, *Escherichia coli*, *Staphylococcus aureus*

**Acknowledgements:** this study was financed by grant ES RTD/2022/7 '3D Biotextile with Technological Composition of Nano Particles to Enhance the Protecting Properties' (3DNano-HPC).

---

## P15: EVALUATION OF RUST AND POWDERY MILDEW DEVELOPMENT IN *MENTHA* × *PIPERITA* AND *MENTHA SPICATA* UNDER *EX SITU* CONDITION

Irina, Sivicka<sup>1</sup>\*, Olga, Sokolova<sup>2</sup>

<sup>1</sup> Institute of Soil and Plant Sciences, Latvia University of Life Sciences and Technologies, latvia

<sup>2</sup> Unit Of Plant Pathology and Entomology, Institute of Horticulture, Latvia

\* Corresponding author. E-mail: Irina.Sivicka@lbtu.lv

---

Mint (*Mentha*), which belongs to the Lamiaceae family, is a widely cultivated aromatic and medicinal plant. However, mint production faces significant threats from fungal diseases. Mint rust, the causal agent *Puccinia menthae*, as well as *Erysiphe cichoracearum*, responsible for powdery mildew, negatively affect both the quantity and quality of raw material. The aim of the work was to evaluate the presence of symptoms of rust and powdery mildew in mint under *ex situ* condition. Study was carried out from 2020 to 2022 using plant material of *Menta* × *piperita* and *Menta spicata* from the *ex situ* collection of genetic resources of aromatic and medicinal plants (N 56°39' 45.3"; E 23°45'15.2") of the Latvia University of Life Sciences and Technologies. During vegetation period, the severity of infection was determined using a scale from 0 to 4, where 0 indicated no disease symptoms and 4 represented 96–100% leaf surface infection. The first symptoms of rust and powdery mildew appeared on most mint accessions in early September. By late October, the severity on plants reached 3–4 points. Differences between *M. × piperita* and *M. spicata*: all *M. × piperita* accessions exhibited symptoms of infection by rust and powdery mildew, ranging from 2 to 4 points, while for *M. spicata* it was from 0 to 4 points. Accession P4 (*M. × piperita*) showed no symptoms of powdery mildew but symptoms of rust infection were reached 2 points. Accessions P9 and P10 (for both *M. × piperita* and *M. spicata*) showed no signs of rust infection but were affected by powdery mildew. The observed differences in development of rust and powdery mildew for *M. × piperita* and *M. spicata* highlight the necessity of studying the genetic basis of resistance of different *Mentha* accessions.

**Keywords:** *Erysiphe cichoracearum*, plant health, *Puccinia menthae*

---

## P16: PHENOLIC COMPOUNDS AND FLAVONOIDS IN MICROALGAL EXTRACTS FOR AGRICULTURAL BIOSTIMULANTS: A COMPARATIVE STUDY OF EXTRACTION METHODS

Tomass Tumpelis\*, Santa Sukaruka, Pāvels Semjonovs

Laboratory of Industrial Microbiology and Food Biotechnology, Institute of Biology, FMLS, University of Latvia

\* Corresponding author. Email: tomass.tumpelis@lu.lv

---

Microalgae *Chlorella vulgaris* and *Arthrospira platensis* are rich in bioactive compounds that are known to contribute to sustainable plant growth. However, variations in extraction methods

significantly influence the concentration of key bioactive compounds, such as phenols and flavonoids, affecting their efficiency as agricultural biostimulants. In this study, varied extraction techniques, i.e., heating, sonication, and microwave-assisted extraction (MAE) were studied to improve the yield of total phenolic compounds as gallic acid equivalent (mg GAE/L) (Anwer et al., 2022) and total flavonoids as quercetin equivalent (mg QE/L) from microalgal biomass into extracts for plant treatment. Results indicated that the highest phenolic and flavonoid content was obtained when biomass (25–200 g/L) was treated by heating at 70°C for one hour, followed by MAE for 120 s using an ethanol-water mixture. Specifically, *A. platensis* with high biomass concentration (200 g/L) yielded approximately 400 mg QE/L flavonoids and 240 mg GAE/L phenols under heating conditions, while *C. vulgaris* yielded 100 mg QE/L flavonoids and 150 mg GAE/L phenols. MAE demonstrated greater efficiency, requiring up to 1/60th of the time than heating to obtain similar phenols and flavonoids extraction yields from microalgal biomass. Under MAE treatment, *A. platensis* flavonoids reached 260 mg QE/L with the lowest biomass concentration (50 g/L). Sonication consistently produced lower phenolic and flavonoid content both in water and water/ethanol (1:1 v/v) mixture, ranging from two to ten times less than under other methods. Comparing efficiency and yield, MAE at 120 s with microalgal biomass concentrations of 100–200 g/L was the optimal extraction technique for both species. These findings highlight the potential of MAE to enhance bioactive compound extraction, providing a more efficient and faster approach for the development of plant biostimulants.

**Keywords:** plant biostimulants, microalgal biomass, microwave-assisted extraction, flavonoids, phenols

**Acknowledgements:** this study was co-financed by European Agricultural Fund for Rural Development (EAFRD) and supported by the Ministry of Agriculture and Rural Support Service of the Republic of Latvia, grant No. 22-00-A01612-0000014.

#### References

1. Anwer SS, Sdiq KH, Muhammad KR, Aladdin LM. Phenolic compound and fatty acid properties of some microalgae species isolated from Erbil City. *Braz J Biol.* 2022;82:e256927.
2. Agustini TW, Suzery M, Sutrisnanto D, Ma'ruf WF, Hadiyanto H. Comparative study of bioactive substances extracted from fresh and dried *Spirulina* sp. *Procedia Environ Sci.* 2015;23:282–289.

---

## P17: RESEARCH ON INSECT BACULOVIRUSES IN LATVIA: PAST AND FUTURE

Līga Jankevica\*

*Department of Experimental Entomology and Microbiology, Institute of Biology, FMLS, University of Latvia, Riga, Latvia*

\* Corresponding author. Email: [liga.jankevica@lu.lv](mailto:liga.jankevica@lu.lv)

---

Research on insect viruses in Latvia began at the Institute of Biology in the last century. The aim was to gain new knowledge on insect viruses and clarify their role in regulating pest populations in Latvia. The main tasks were to monitor significant forest pest populations, to obtain new isolates and describe their properties, and to investigate occurrence and natural variability of baculoviruses in pest populations. After the 1990s, the focus was on the development of experimental strains with high virulence, development of novel viral insecticide preparations, determination



of their efficacy, and assessment of the perspectives for use in pest control. The persistence of viruses in biocenosis (soil, on plant leaves, in water, etc.) was studied. In the early 21st century, investigations on interaction of baculoviruses in the plant-pest-pathogen system and the studies of baculoviruses genome were carried out.

Monitoring of outbreaks and natural epizootics of forest pests have been done on a regular basis. Living insects were laboratory-reared in isolators under optimal conditions. The presence of baculoviruses in larval tissue was detected by direct examination of larval tissue smears (Lacey, Brooks, 1997). The viral polyhedra or granules were observed in an electron microscope. Methods of DNA amplification by specific primers were used for identifying latent and persistent viral infections. Biotests were carried out for determining the virulence and efficacy of isolates. Virulent experimental strains were obtained from wild isolates, using passages through host organisms under stress factors.

The majority of entomopathogenic viruses were isolated from dangerous forest and agricultural pests. Most of the isolated and identified insect virus isolates belong to the family Baculoviridae, 13 of them belong to the genus *Alphabaculoviruses*, six to the genus *Betabaculoviruses*, and four to the genus *Gammabaculoviruses*, a viral insecticide VIRIN KS against the European tent caterpillar or lackey moth *Malacosoma Neustria* L. and VIRIN NS for use against the pine sawfly *Neodiprion sertifer* was developed on the basis of experimental strains in the last century (Zariņš, Eglite, 1993). Methods of developing dry and liquid virus insecticide forms were patented. Based on the polyhedrin gene sequence (GenBank AJ277555.1), specific primers were created for the identification of MnMNPV in the environment and pest population. Observations showed that declining populations of *Malacosoma neustria* and *Acantholyda posticalis* had latent or sublethal NPV infection (Jankevica et al., 2002; Jankevica et al., 2020). In recent years, the role of microorganism associations of the middle intestinal microflora in development of baculovirus infection was clarified in collaboration with JKI (Germany) and ETH Zurich (Swiss).

**Keywords:** Baculoviridae, biological control, persistent viral infections, virus insecticide

#### References

1. Lacey LA, Brooks WM. Initial handling and diagnosis of diseased insects. In: Manual of techniques in insect pathology. Lacey, LA, editor. London: Academic Press; 1997. p. 1–15.
  2. Jankevica L, Kropa M, Savenkovs N, Jankevics E. Presence of Nucleopolyhedroviruses in Natural Populations of *Malacosoma neustria* L. (Lepidoptera, Lasiocampidae) Latvijas Entomologs, 2002;39:30–7.
  3. Jankevica L, Striķe Z, Smits A. Preliminary studies on presence of entomopathogens in outbreak population populations of the great web spinning pine sawfly *Acantholyda posticalis* in urban forest stands. IOBC-WPRS Bulletin. 2020;150:161–4.
  4. Zariņš I, Eglite G. Investigation of entomopathogenous viruses in Latvia and their potential as pest control. Proc Latv Acad Sci. 1993;557:49–53.
-